

Nonfactorizable effects in $B \rightarrow J/\psi K$

Blaženka Melić^{1,2} * ** and Reinhold Rückl¹

¹ Institut für Theoretische Physik und Astrophysik, Universität Würzburg, D-97074 Würzburg, Germany

² Institut für Physik, Universität Mainz, D-55099 Mainz, Germany

Abstract. The method of QCD light-cone sum rules is used to calculate soft nonfactorizable contributions to the decay amplitude for $B \rightarrow J/\psi K$. The result confirms expectations that in color-suppressed decays nonfactorizable corrections can be sizable.

1 Decay amplitude

Precise measurements of exclusive nonleptonic B decays have initiated theoretical considerations which go beyond *naive factorization* frequently used in the calculation of decay amplitudes. Nonfactorizable contributions have been investigated in several approaches [1,2,3] for different classes of two-body nonleptonic B decays. Here, we focus on nonfactorizable corrections in the decay $B \rightarrow J/\psi K$. This channel is particularly interesting because of a substantial discrepancy between the prediction from naive factorization and experiment by more than a factor of three in the branching ratio. Also, this mode belongs to the color-suppressed class-2 decays for which large nonfactorizable contributions are expected.

The weak matrix element can be written in the form

$$\langle J/\psi K | H_W | B \rangle = \sqrt{2} G_F V_{cb} V_{cs}^* \epsilon \cdot q m_{J/\psi} f_{J/\psi} F_{BK}^+(m_{J/\psi}^2) a_2, \quad (1)$$

where F_{BK}^+ is the $B \rightarrow K$ form factor and the parameter a_2 incorporates factorizable and nonfactorizable contributions. A particular useful parametrization is given by

$$a_2 = C_2(\mu) + \frac{C_1(\mu)}{N_c} + 2 C_1(\mu) \frac{\tilde{F}_{BK}^+(\mu)}{F_{BK}^+(m_{J/\psi}^2)}, \quad (2)$$

where $C_{1,2}$ are the short-distance Wilson coefficients. The first two terms in (2) result from naive factorization, while the term proportional to $\tilde{F}_{BK}^+$ represents nonfactorizable contributions. Since a_2 parametrizes a physical decay amplitude, the scale dependence of the individual terms in a_2 must cancel. Taking $\mu = m_b$, one has, numerically, $C_1(m_b) = 1.802$ and $C_2(m_b) = -0.185$ [4].

* Talk given by B.M. at the Workshop on B physics and CP violation, Frontier Science 2002, 06-11 October 2002, INFN, Frascati, Italy, to appear in the Frascati Physics Series.

** Alexander von Humboldt fellow. On leave of absence from Rudjer Bošković Institute, Zagreb, Croatia.




Fig. 1. Leading soft nonfactorizable contributions as estimated by *QCD light-cone sum rules*. The shaded ellipse denotes the *K* meson light-cone distribution amplitude. The currents $j_\nu^{J/\psi}$ and j_5^B generate states with *J/ψ* and *B* quantum numbers, respectively. The square stands for the $\tilde{O}_2$ four-quark weak operator [6].

2 Nonfactorizable effects

According to the large N_c analysis [5], there could be a cancellation between the last two terms in (2), both being of order $1/N_c$ relative to the leading term. In case of such a cancellation, $a_2 \simeq C_2$ would be negative. While for *D* decays a negative value of a_2 is consistent with experiment, experimental data on *B* decays suggest a positive a_2 . On the other hand, explicit estimates of nonfactorizable contributions in *B* decays have predicted both signs for a_2 .

In [6], the *QCD light-cone sum rule method* [3] was applied to the calculation of soft nonfactorizable contributions in $B \rightarrow J/\psi K$. The relevant operators in the weak Hamiltonian H_W are

$$\mathcal{O}_2 = (\bar{c}\Gamma_\mu c)(\bar{s}\Gamma^\mu b), \quad \tilde{\mathcal{O}}_2 = (\bar{c}\Gamma_\mu \frac{\lambda^a}{2} c)(\bar{s}\Gamma^\mu \frac{\lambda^a}{2} b). \quad (3)$$

It is the contribution of $\tilde{\mathcal{O}}_2$ to the matrix element (1), which is expected to give rise to the leading nonfactorizable effects. The sum rule approach allows to estimate the contribution of soft-gluon exchange between the *J/ψ* and the *B*–*K* system, see Fig. 1. To this end, one has to isolate the ground state contribution to the correlation function

$$F_\nu(p, q, k) = \int d^4x e^{-i(p+q)x} \int d^4y e^{i(p-k)y} \langle K(q) | T \{ j_\nu^{J/\psi}(y) \tilde{\mathcal{O}}_2(0) j_5^B(x) \} | 0 \rangle \quad (4)$$

represented graphically in Fig.1. Taking into account twist-3 and twist-4 contributions calculated at the appropriate scale $\mu_b \sim m_b/2$, one finds

$$\tilde{F}_{BK}^+(\mu_b) = 0.009 \div 0.017, \quad (5)$$

and, substituting (5) in (2),

$$a_2 = 0.14 \div 0.17|_{\mu=\mu_b}. \quad (6)$$

Although the nonfactorizable matrix element (5) is rather small, it increases a_2 by $30 \div 70\%$, due to the large coefficient $2C_1(\mu_b)$.

In addition, one has nonfactorizable contributions from hard-gluon exchange. They have been estimated in *QCD factorization* [1] and amount to another 20% correction [7]. Thus, in total one obtains

$$a_2 = 0.16 \div 0.19|_{\mu=\mu_b}. \quad (7)$$

3 Conclusions

Firstly, one can see that the theoretical expectation (7) is still too small to explain the experimental value

$$|a_2|^{exp} = 0.29 \pm 0.03. \quad (8)$$

Secondly, it is interesting to note that the theoretical approach described here predicts a positive sign of a_2 in agreement with experiment and in contradiction with the argument based on $1/N_c$ expansion [5]. Finally, a comparison of the value (8) deduced from the measurement of $B \rightarrow J/\psi K$ with the result $|a_2|^{exp} = 0.4 \div 0.5$ from the decays $\bar{B}^0 \rightarrow D^{(*)0}\pi^0$ indicates a substantial nonuniversality of a_2 in color-suppressed decays [8].

Acknowledgments

B.M. wishes to thank the organizers of Frontier Science 2002 for the invitation and for their support and would also like to acknowledge the support by the Alexander von Humboldt Foundation and partial support of the Ministry of Science and Technology of the Republic of Croatia under the contract 0098002. R.R. acknowledges the support by the Bundesministerium für Bildung und Forschung (BMBF, Bonn, Germany) under the contract number 05HT1WWA2.

References

1. M. Beneke, G. Buchalla, M. Neubert, C.T. Sachrajda, Nucl. Phys. B **591**, 313 (2000); Nucl. Phys. B **606**, 245 (2001).
2. Y.-Y. Keum, H.-n. Li, A.I. Sanda, Phys. Lett.B **504**, 6 (2001); Phys. Rev. D **63**, 054008 (2001).
3. A. Khodjamirian, Nucl. Phys. B **605**, 558 (2001).
4. G. Buchalla, A.J. Buras, M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
5. A.J. Buras, J.M. Gerard, R. Rückl, Nucl. Phys. B **268**, 16 (1986).
6. B. Melić, R. Rückl, Soft nonfactorizable corrections in $B \rightarrow J/\psi K$, preprint WUE-ITP-2002-020, MZ-TH-02-23, in preparation.
7. H.-Y. Cheng, K.-Ch. Yang, Phys. Rev. D **63**, 074011 (2001).
8. M. Neubert, A.A. Petrov, Phys. Lett. B **519** (2001) 50.